

MP Biomedicals, LLC

29525 Fountain Parkway Solon, Ohio 44139 Telephone: 440/337-1200 Toll Free: 800/854-0530 Fax: 440/337-1180 mailto: biotech@mpbio.com

mailto: <u>biotech@mpbio.com</u> web: <u>http://www.mpbio.com</u>

TECHNICAL INFORMATION

Catalog Number: 960438, 960439, 960440, 960441

Methionine/Choline Deficient and Control Diets

Description: Nonalcoholic steatohepatitis (NASH) is common liver injury in which the histopathological abnormalities mimic those of alcoholic steatohepatitis. This disorder is most frequently associated with type 2 diabetes, hyperlipidemia, and rapid weight loss. Leclercq, et al. published in The Journal of Clinical Investigation, CYP2E1 and CYP4A as microsomal catalysts of lipid peroxides in murine nonalcoholic steatohepatitis, in which they proposed that CYP2E1 could also be a factor in the development of NASH. The mice used in this experiment were fed MP's plain Methionine/Choline and our control diet.

References:

– Leclercq, I.A. (Storr Liver Unit, Department of Medicine and Westmead Millennium Institure, University of Sydney at Westmead Hospital, Westmead NSW 2145, Australia), Farrell, G.C. (Storr Liver Unit, Department of Medicine and Westmead Millennium Institure, University of Sydney at Westmead Hospital, Westmead NSW 2145, Australia), Field, J. (Storr Liver Unit, Department of Medicine and Westmead Millennium Institure, University of Sydney at Westmead Hospital, Westmead NSW 2145, Australia), Bell, D.R. (School of Biology, University of Nottingham, University Park, Nottinhgam NG7 2DR, United Kingdom), Gonzalez, F.J. (Laboratory of Molecular Carcinogenesis, National Cancer Institute, National Institutes of Health, Bethesda, Maryland 20892, USA) and Robertson, G.R. (Storr Liver Unit, Department of Medicine and Westmead Millennium Institure, University of Sydney at Westmead Hospital, Westmead NSW 2145, Australia), "CYP2E1 and CYP4A as microsomal catalysts of lipid peroxides in murine nonalcoholic steatohepatitis." *The Journal of Clinical Investigation*, v. 10(8), 1067-1075.

Methionine/Choline Deficient Diet Composition:

Ingredient Sucrose Corn Starch Corn Oil Alphacel Non-Nutritive Bulk AIN 76 Mineral Mix (composition below)	Amount (g/kg diet) 455.3 gm 200.0 gm 100.0 gm 30.0 gm 35.0 gm
	(below is gm/kg of AIN 76 Mineral Mix)
Calcium Phosphate Dibasic	500.00 gm
Sodium Chloride	74.00 gm
Potassium Citrate Monohydrate	220.00 gm
Potassium Sulfate	52.00 gm

Magnesium Oxide	24.00 gm
Manganese Carbonate (43-48% Mn)	3.50 gm
Ferric Citrate (16-17% Fe)	6.00 gm
Zinc Carbonate (70% ZnO)	1.60 gm
Cupric Carbonate (53-55% Cu)	0.30 gm
Potassium Iodate	0.01 gm
Sodium Selenite	0.01 gm
Chromium Potassium Sulfate	0.55 gm
Sucrose, finely powdered	118.00 gm
, , , , , , , , , , , , , , , , , , ,	· ·
Dicalcium Phosphate L-Alanine L-Arginine Hydrochloride L-Asparagine Monohydrate L-Aspartic Acid L-Cystine L-Glutamic Acid Glycine L-Histidine Hydrochloride L-Isoleucine L-Leucine L-Leucine L-Phenylalanine L-Proline L-Serine L-Threonine L-Tryptophan L-Tyrosine L-Valine DL-alpha-Tocopherol Acetate (250 u/gm) Vitamin A Palmitate (250,000 u/gm) Ethoxyquin	3.0 gm 3.5 gm 12.1 gm 6.0 gm 3.5 gm 40.0 gm 23.3 gm 4.5 gm 8.2 gm 11.1 gm 18.0 gm 7.5 gm 3.5 gm 3.5 gm 8.2 gm 1.8 gm 5.0 gm 8.2 gm 0.484 gm 0.0792 gm 0.0055 gm 0.002 gm
Dicalcium Phosphate L-Alanine L-Arginine Hydrochloride L-Asparagine Monohydrate L-Aspartic Acid L-Cystine L-Glutamic Acid Glycine L-Histidine Hydrochloride L-Isoleucine L-Leucine L-Leucine L-Phenylalanine L-Proline L-Serine L-Threonine L-Tryptophan L-Tryrosine L-Valine DL-alpha-Tocopherol Acetate (250 u/gm) Vitamin A Palmitate (250,000 u/gm) Vitamin D3 (400,000 u/gm)	3.0 gm 3.5 gm 12.1 gm 6.0 gm 3.5 gm 40.0 gm 23.3 gm 4.5 gm 8.2 gm 11.1 gm 18.0 gm 7.5 gm 3.5 gm 3.5 gm 8.2 gm 1.8 gm 5.0 gm 8.2 gm 0.484 gm 0.0792 gm 0.0055 gm 0.002 gm
Dicalcium Phosphate L-Alanine L-Arginine Hydrochloride L-Asparagine Monohydrate L-Aspartic Acid L-Cystine L-Glutamic Acid Glycine L-Histidine Hydrochloride L-Isoleucine L-Leucine L-Lysine Hydrochloride L-Phenylalanine L-Proline L-Serine L-Threonine L-Tryptophan L-Tyrosine L-Valine DL-alpha-Tocopherol Acetate (250 u/gm) Vitamin A Palmitate (250,000 u/gm) Ethoxyquin	3.0 gm 3.5 gm 12.1 gm 6.0 gm 3.5 gm 40.0 gm 23.3 gm 4.5 gm 8.2 gm 11.1 gm 18.0 gm 7.5 gm 3.5 gm 3.5 gm 8.2 gm 1.8 gm 5.0 gm 8.2 gm 0.484 gm 0.0792 gm 0.0055 gm 0.002 gm
Dicalcium Phosphate L-Alanine L-Arginine Hydrochloride L-Asparagine Monohydrate L-Aspartic Acid L-Cystine L-Glutamic Acid Glycine L-Histidine Hydrochloride L-Isoleucine L-Leucine L-Lysine Hydrochloride L-Phenylalanine L-Proline L-Serine L-Threonine L-Trytophan L-Tyrosine L-Valine DL-alpha-Tocopherol Acetate (250 u/gm) Vitamin A Palmitate (250,000 u/gm) Ethoxyquin Vitamin Mix	3.0 gm 3.5 gm 12.1 gm 6.0 gm 3.5 gm 40.0 gm 23.3 gm 4.5 gm 8.2 gm 11.1 gm 18.0 gm 7.5 gm 3.5 gm 3.5 gm 8.2 gm 1.8 gm 5.0 gm 8.2 gm 0.484 gm 0.0792 gm 0.0055 gm 0.002 gm

Inositol	0.1101 gm
Menadione	0.0496 gm
	0.0991 gm
Niacin	0.1101 gm
p-Aminobenzoic Acid	0.0220 gm
Pyridoxine Hydrochloride	0.022 gm
Riboflavin	0.022 gm
Thiamine Hydrochloride	0.0297 gm
Vitamin B12 (0.1% trit.)	1.0166 gm
Ascorbic acid	3.4503 gm
Cara Starah	

Methionine/Choline Control Diet Composition:

This diet has the same formulation as the Methionine/Choline Deficient diet except we add 2 g/kg choline chloride and 3 g/kg DL-Methionine at the expense of sucrose.

Availability:

Corn Starch

Catalog Number	Description	Sizes
		10 kg 20 kg 50 kg
		10 kg 20 kg 50 kg
		10 kg 20 kg 50 kg
960441		10 kg 20 kg 50 kg